

Robert Ullman, N.D. and Judyth Reichenberg-Ullman, N.D.
The Northwest Center for Homeopathic Medicine
123 4th Avenue, N., Suite 2 Edmonds, WA 98020 (360) 221-7715
nchmclinic@gmail.com

How to Take Your Liquid Homeopathic Medicine

Plussed Medicines:

- You will receive a 1-oz. dropper bottle containing your liquid remedy. You will use the same bottle until it is empty or we instruct you to stop.
- Shake the bottle against your hand or a book 10 times.
- Place 5 drops under the tongue with the frequency directed. (For children, you may be advised to put the 5 drops in an oz. or so of water and sip to avoid the alcohol taste.)
- If you do not want any alcohol, just let us know and you will then need to refrigerate the bottle to preserve it.
- (If Dr. Ullman directs you to dilute the liquid further, put the 5 drops in 4 oz. of water, stir, and take ½ tsp. You will make it freshly each time you take it.)

LM Medicines:

- You will receive a 1-oz dropper bottle containing your remedy. You will be making a new dilution each day you are instructed to take the medicine.
- Shake the bottle against your hand or a book 10 times.
- Place 5 drops in 4 oz. water in a paper cup.
- Stir this solution for 15 seconds vigorously with a spoon.
- Take ½ tsp. of the stirred solution.
- Throw away the remaining solution.

LM Medicines Multiple-Glass Method for Sensitive Individuals:

- Follow the instructions above to prepare the first glass of the LM medicine.
- Instead of taking the ½ tsp. of the liquid, put it in a second 4-oz glass of water. Stir that glass and take ½ tsp. of it.
- You may be asked to repeat this process a specific number of additional times if you are very sensitive.